PREVENTION OF PRESSURE ULCERS





ABOUT PRESSURE ULCERS

A pressure ulcer is a localized injury of the skin and/or underlying tissue usually over a bony prominence, caused by pressure, or combination of pressure and shear.



1 to 4 out of 10

hospitalized patients have pressure ulcers(1)



50%

of pressure ulcers are preventable, depending on population and ulcer grade (2)

HOW TO PREVENT PRESSURE ULCERS(3)?



Healthy body weight & Nutrition



Quit Smoking



Hydration and moisture



Skin inspection



Ease pressure and prevent friction



Spread awareness

AREAS AT RISK







WHO IS AT RISK OF PRESSURE ULCERATION?



Reduced mobility/activity



History of pressure ulceration



Impaired circulation due to diabetes, vascular problems, oedema




Increased skin humidity (eg due to incontinence, perspiration)



Poor nutritional status

B. BRAUN SOLUTIONS





Hydrophilic foam dressings Anatomically shaped hydrocellular heel dressing



Askina® DresSil® Sacrum

Foam dressings with silicone adhesive



Linovera®

Solution of hyperoxygenated fatty acids
In a prospective study, none of the high

In a prospective study, none of the high risk patients preventively treated with Linovera® developed pressure injuries. (4)



Askina® Barrier Cream

Protectant and moisture barrier

This document, its contents, including institutional data, information, trademarks and logos mentioned herein are the exclusive property of B. Braun. Any representation and/or reproduction, total or partial, of this document and its contents without the express prior consent of B. Braun, is strictly prohibited and constitutes an infringement of the intellectual property rights of B. Braun.

Non-binding documents and photographs. For healthcare professional use only.

^{1.} Range varies among settings and classification method. https://www.bbraun.com/en/products-and-therapies/wound-management/prevention-of-pressure-ulcers.html 2. Downie, F. & Sandoz, Heidi & Gilroy, P. & Royall, Dawn & Davies, S.. (2013). Are 95 % of hospital-acquired pressure ulcers avoidable? Wounds UK. 9. 16-22.

^{3.} http://www.internationalguideline.com/static/pdfs/NPUAP-EPUAP-PPIA-PUQuickReferenceGuide-2016update.pdf

^{4.} López-Pérez, R. Gutiérrez Ibáñez, B. Incidence of pressure ulcers (bed sores) in patients on treatment with hyperoxygenated fatty acids. (Asturias, Spain). Unpublished.

B. Braun Hospicare Ltd. | Collooney | Co. Sligo | Ireland
B. Braun Medical SA | Carretera de Terrassa 121 | 08191 Rubi (Barcelon)